

## Violence Crisis Services During Lockdown: An Assessment of Service Availability During COVID-19 Lockdown in South Africa

### Introduction

#### *The state of gender-based violence in South Africa*

Prior to COVID-19, South Africa faced a well-documented epidemic of gender-based violence (GBV). The most recent South African Demographic and Health Survey (DHS) reports that more than a quarter (26%) of ever-partnered South African women have experienced any type of physical, sexual, or emotional violence by a partner. The DHS also reported that the proportion of women reporting violence by a partner in the last year varied significantly by province – as low as 7% in Limpopo and as high as 18% in North West.<sup>1</sup>

The South Africa GBV crisis has received attention at the international and national level, prompting government and civil society responses. The recent R1.6 billion Emergency Response Action Plan on Gender-based Violence and Femicide represents the latest example of government efforts to invest in expanding GBV service access.<sup>2</sup> Nongovernmental organizations (NGOs) also play a significant role in the GBV response in terms of advocacy, prevention, and response.<sup>3</sup> Accordingly, GBV services in South Africa are diverse in terms of both the entities providing them and available services.

For the purpose of this analysis, we focus specifically on two types of services: Thuthuzela Care Centers (TCCs) and domestic violence shelters, given their importance in the immediate response to GBV survivors. However, we acknowledge this is not the sum of GBV services in South Africa, and in fact, globally, many GBV survivors rely on informal social support rather than accessing formal services.<sup>4,5</sup>

TCCs are one-stop sexual assault centers that aim both to aid in conviction of sexual offenses but also to expand physical, psychological and social care for survivors of rape, sexual assault and domestic violence.<sup>6</sup> While the primary route through which individuals access TCCs is via police referrals and transport, TCCs also receive referrals from hospitals and other health care

### Key Findings

- Government and civil society have made significant efforts to keep gender-based violence (GBV) services available during the COVID-19 lockdown, a time that may exacerbate challenges for those at risk of GBV.
- Despite this commitment, a rapid assessment of violence services in South Africa finds that a quarter of Thuthuzela Care Centers – government-run, one-stop sexual assault resource centers – and 40% of violence shelters were unreachable during a week of twice-daily calling.
- Additionally, only 25 shelters nationwide stated they were able to take in new clients, and not all individuals seeking services would likely meet the criteria for admission.
- These troubling barriers to accessibility are likely indicative of COVID-19 lockdown challenges as well as existing pre-COVID-19 deficiencies in South African GBV service capacity.
- Further government action and resources must be directed to ensure reliable access to GBV services for the duration of COVID-19 and beyond.

providers, as well as individuals coming by themselves directly to the centers.<sup>7</sup> While few peer-reviewed evaluations of TCCs exist, the grey literature suggests that individuals seeking care at TCCs have faced documented delays in care and other issues including inadequate privacy.<sup>8,9</sup> Critiques of TCCs have also found a lack of resources for sufficient follow-up and comprehensive service delivery, noting that while acute medical needs may be met, emotional support services are often not provided due to insufficient funds, space, and social workers.<sup>10</sup>

Violence shelters are another vital aspect of the GBV response, including the National Shelter Movement of South Africa (NSMSA), which serves as a collective voice for the majority of South Africa's GBV shelters.<sup>11</sup> Similar to TCCs, the grey literature suggests there are insufficient violence shelters in South Africa, many of which have overburdened staff serving in multiple roles, and rely heavily on volunteers. Many shelters have had to cut program provision as expenditures exceed Department of Social Development funding.<sup>12</sup>

In addition to limited capacity, needs also may not be met for some survivors of GBV because of various criteria required for being housed within a shelter. A review of these criteria includes examples such as some shelters barring women with children, women with male children over a certain age, individuals with mental health challenges, and/or men.<sup>13</sup> Attention has also been called to the specific challenges individuals with disabilities face in accessing GBV services, including lack of accommodations for physical disabilities and communication challenges.<sup>14</sup> Additionally, shelters may not be able to provide appropriate resources to survivors of same-sex intimate partner violence.<sup>15</sup> In short, challenges to finding GBV services existed for many in South Africa, even prior to the COVID-19 crisis.

### There are insufficient violence shelters in South Africa, many of which have overburdened staff serving in multiple roles.

#### *Gender-based violence and COVID-19*

COVID-19-related lockdowns and layoffs can increase risk of GBV by simultaneously isolating people with their violent partners during times of economic and social stress and cutting them off from available resources.<sup>16</sup> Past global health crises have demonstrated these risks; quarantines and school closures during the 2014 Ebola outbreak resulted in documented harms, including sexual coercion and abuse, to young women.<sup>17</sup> Moreover, in addition to direct harms sustained by individuals during such public health crises, GBV service delivery is often disrupted when personnel and resources are redirected for emergency response.<sup>18</sup>

Cognizant of the lockdown's potential exacerbation of the existing GBV crisis, the South African government and civil society have moved quickly to adapt GBV resources to better serve survivors during these unprecedented times. In particular, measures to mobilize virtual resources have been vital to the response, including bolstering 24-hour hotlines for domestic violence and rape crisis, and providing virtual support via Skype,

"Please call me" SMS message services, telephone counseling, and other safety-planning services and shelter referrals for survivors.<sup>19,20</sup> Additionally, the government classified TCCs and shelter services as essential services during the COVID-19 pandemic to allow continued provision of care.<sup>21,22</sup>

In the initial period after lockdown began, the national GBV Command Center saw a small increase in calls, but other GBV-related services, including cases reported to TCCs, decreased.<sup>21</sup> Decreased reporting is likely not indicative of decreases in GBV, but often reflective of decreased ability to access services,<sup>23</sup> further emphasizing the importance of ensuring access to services during lockdown. This rapid assessment aims to assess the extent to which these GBV services remained operational during COVID-related lockdowns in South Africa.

### Methods

A rapid assessment of GBV crisis services was conducted by amfAR, The Foundation for AIDS Research, to record whether services remained open and providers were answering the phone during COVID-related lockdowns. Data collection spanned one month (April 30–May 29) during which South Africa was in lockdown level 5 (enforced restrictions on all non-essential movement) or level 4 (slightly eased regulations on movement with strict restrictions on most travel).<sup>24</sup> During the assessment, a team of six data collectors called all publicly listed gender-based violence shelters, using the primary phone number listed for each shelter in the National Shelter Movement of South Africa (NSMSA) shelter directory (n=86)<sup>25</sup> and all TCCs listed by the National Prosecuting Authority of South Africa (n=55).<sup>26</sup> Seven additional known shelters through researcher contacts were added to the initial list (two shelters in Gauteng province, one in KwaZulu Natal, and two in Mpumalanga province). Total shelter count n=93.

Data collectors called every GBV service twice a day until a person was reached for up to seven days, or a maximum of 14 calls per service.\* Calls were all made during South African business hours and from a local South African phone number. Callers identified themselves as researchers based in the United States when asked, otherwise the questions were asked without the researchers providing additional background information on themselves. Of primary interest were whether publicly available numbers for GBV crisis services were working and whether these services continued to field incoming calls during COVID lockdowns.

GBV shelters with intake capacity were asked if they had space available to house new clients. If staff described any criteria for admittance to the shelter, that was also noted but was not

\* Five of the 19 shelters that were deemed non-responders were only contacted 4 times instead of the full 14 calls due to time constraints.

specifically asked by data collectors. Survivors who are seeking shelter may not always be contacting shelters directly, but rather through referral from someone who may help them navigate the shelter system such as a social worker, police, NSMSA representative, or hotline, etc. This is particularly true given the specific referral system set up for shelters during COVID-19.<sup>19,20</sup>

However, the decision was made to call the main numbers for each shelter listed in the NSMSA directory<sup>25</sup> for the following reasons. First, the study team sought information specific to each shelter, and second, with the directory remaining prominently listed on the NSMSA website, it was assumed this was a place survivors seeking support might still visit.

It is probable that some GBV survivors seeking care may have been able to find the appropriate instructions to reach the shelter representative for each province. Thus this data collection effort is not meant to exactly replicate a survivor’s search for services but instead to capture information regarding the state of services during COVID-19 and note the challenges some might face in reaching a shelter. The difficulties that data collectors experienced in trying to find current publicly available contact information for crisis may be faced by survivors as well.

Other limitations to this assessment are that TCCs and shelters do not make up the totality of South Africa’s GBV response, and data collectors did not attempt to contact hotlines or social workers specifically. For ethical reasons the study team chose not to add call volumes to hotlines or individuals involved in case

**The difficulties that data collectors experienced in trying to find current publicly available contact information for crisis may be faced by survivors as well.**

management, and therefore this assessment does not reach the totality of the GBV response system in South Africa. Accordingly, the results should be interpreted only as an initial indication of service availability through the method of contact we selected. Further investigation is needed to assess the accessibility of GBV services in the unique setting of the COVID-19 lockdowns.


**Results**

Overall, the assessment found that 63% of TCCs and shelters were reachable during the assessment, requiring an average of three calls per service. Of the known and reachable GBV shelters, 25 confirmed that they had space to accommodate new clients for intake [Figure 1].

*Thuthuzela Care Centers (TCCs)*

TCCs are located in each province, with four each in Free State and Northern Cape, five in Mpumalanga and North West, seven each in Gauteng, Limpopo, and Western Cape, and eight in KwaZulu-Natal and Eastern Cape. During the data collection process, two TCCs of the original 55 were determined to have

**Figure 1. GBV services by type and availability, May 2020\***



\* Exact shelter location unavailable for safety reasons. Locations represent approximations based on available information in the National Shelter Movement directory.

been decommissioned. Of the remaining 53, the majority (40, 75%) were reachable by phone during the week of calling. Thirteen TCCs were not reachable by phone. Twenty-two TCCs answered the phone on the first attempt. On average, it required 2.5 calls to a facility to reach a live person (range 1–12 calls) [Figure 2]. In general, data collectors found TCC staff to be friendly and informative when reached. Many TCCs had phone numbers that routed through the larger hospital facility number, and occasionally main hospital staff were unsure about the existence of TCCs. Given that TCC services rely significantly on referrals from police rather than individuals seeking the services by themselves, it is not unexpected that directly calling the TCCs was sometimes a challenge. However, in the event that a survivor or patient reaches the TCC directly, phone calls should still be fielded accordingly.

Of the reachable TCCs, 26 (65%) confirmed that they were open 24 hours a day to serve clients – although not all services would be available at all times. Several TCCs noted that while services were not available 24 hours a day on-site, they partnered with other NGOs that could provide more specialized services during non-business hours like nights, weekends, and holidays. These TCCs were still counted as operating 24 hours a day.

#### Shelters

The number of shelters varied by province: one in North West, two in Limpopo and Northern Cape, four in Free State, seven


in Eastern Cape, nine Kwa-Zulu Natal, 16 in Mpumlanga, 18 in Western Cape and 28 in Gauteng province. Nearly a quarter (23%) of the shelters' listed telephone numbers were nonfunctional (i.e., number was immediately disconnected or call could not be completed). Among unreachable shelters, data collectors were not able to distinguish between those with outdated phone numbers, those that may have no longer been functioning prior to COVID-19, and those closed due to COVID-related issues.

Fifty-three shelters (57%) were ultimately reachable during the assessment, and 35 shelters answered on the first attempt. Of the shelters with working numbers, it took data collectors an average of four calls to reach a live person (range 2–7 calls) [Figure 3]. At the time of data collection, 25 shelters across the country reported having room for new clients who met the shelter's client criteria. Similar to the TCCs, data collectors found that when they were able to reach staff, they were friendly and informative; however, few shelter staff were able to confirm that they could accommodate new clients for intake, though some were able to provide the contact information of social workers with that information.


The specific criteria required for shelter entry varied. Most shelters required a client to be a woman and any accompanying children to be under a certain age. Entry into one shelter in Gauteng was contingent on proof of a negative COVID-19 test and another in Eastern Cape required clients to provide their own personal protective equipment – requirements that could make the shelter inaccessible for many. Two shelters indicated they could only accept intake clients with a referral from the Department of Social Development (DSD) but were open 24 hours a day given their government designation as first responders. Both of these shelters were located in Western Cape, where a specific two-tier referral system had been put into place in which several shelters were identified as Stage One, where survivors spend two weeks to monitor for COVID symptoms before moving on to Stage Two shelters.<sup>27</sup>

This assessment documented only a small snapshot of the barriers that a survivor may need to overcome to access a shelter. Additional barriers may include lack of transportation, financial constraints, citizenship requirements, or restrictions on the number and/or age of accompanying children. Together, these barriers have the potential to exclude survivors from shelter even if there is space available for intake.

**Figure 2. TCC Accessibility**



**Figure 3. Shelter Accessibility**



individuals who made hundreds of calls to service providers across South Africa during this rapid assessment, an actual survivor may live in a province with only one shelter and only have time to make one call. A non-answered call or incorrect published phone number should not join the countless other social, cultural, and economic barriers that survivors of violence need to overcome in order to secure their safety.

COVID infection does pose a serious risk to clients and staff of GBV services that cannot be ignored. Many shelters in this assessment have taken steps to reduce COVID risk for their clients and staff such as requiring a negative COVID-19 test prior to admittance, requiring individuals to bring their own PPE, reducing shelter capacity, or the two-tier system implemented in Western Cape. Given that some of these requirements may be prohibitive for those seeking services, government support to ensure availability and accessibility of needed COVID-19 testing or PPE is vital for the GBV response in order to keep both clients and staff safe.

Even as lockdowns ease, COVID-19-related food insecurity and economic instability, both of which have been associated with GBV,<sup>30-31</sup> will remain prevalent. A significant increase in governmental and donor resources for social workers, case management, intake services,

and active outreach and dissemination of GBV information will be required to combat the GBV pandemic in the face of COVID-19. In President Ramaphosa’s May 13th (2020) speech, he stressed that lockdown measures were designed to ensure that survivors of violence would continue to receive assistance.<sup>32</sup> This promise needs to be backed up by the resources necessary to ensure that a survivor’s call for help is answered the first time, every time.

Of note, the assessment also found that shelters were adapting to the challenges of COVID-19 by providing additional virtual support, phone-based counseling, and safety planning. Some shelters were maintaining limited vital functions, such as handing out food and clothing, despite the inability to accommodate new clients for intake. Those that were not accepting new clients during lockdown planned to be fully operational after lockdowns eased, though the actual ability of these shelters to accommodate new clients post-lockdown is not guaranteed. Indeed, many of the challenges noted during this rapid assessment, including limited shelter space and staff capacity to field calls, may be indicative of pre-existing issues that will not be addressed as lockdowns ease.

**Conclusions**

While there have been noteworthy adaptations to service delivery by South African GBV service providers during COVID-19, this assessment finds that actual availability of services during lockdown may be more limited than what is publicly listed due to non-working numbers and non-answered phone calls during these difficult times. Unlike the well-resourced, non-traumatized

**Acknowledgments**

Thank you to all of the dedicated South African service providers who took the time to respond to our questions and for their continued work to provide GBV services during the COVID-19 crisis. Thank you to the amfAR staff who made this project possible, including Jennifer Sherwood, Elise Lankiewicz and Brian Honermann, and to the hardworking data collectors who volunteered their time to make this project a reality, including Brooke Wurst, Cara Guenther, Elizabeth Saltonstall, and Nozipho Zungu.

## REFERENCES

1. National Department of Health (NDoH), Statistics South Africa (Stats SA), South African Medical Research Council (SAMRC), and ICF. South Africa Demographic and Health Survey 2016. Pretoria: National Department of Health, 2019. <https://dhsprogram.com/pubs/pdf/FR337/FR337.pdf> (accessed 14 July 2020).
2. Interim Steering Committee on Gender-Based Violence and Femicide. Emergency Response Action Plan on Gender-Based Violence and Femicide. 2020. <http://www.thepresidency.gov.za/download/file/fid/1853> (accessed 28 May 2020).
3. Centre for the Study of Violence and Reconciliation. Mapping local gender-based violence prevention and response strategies in South Africa. Braamfontein. <https://www.csvr.org.za/pdf/Mapping-gender-based-violence-prevention-and-response-strategies.pdf> (accessed 14 July 2020).
4. Klein, R. Sickening Relationships: Gender-Based Violence, Women's Health, and the Role of Informal Third Parties. *Journal of Social and Personal Relationships*. 2004;(21)1:149-165.
5. Rasool, S. Do We Accept the Unacceptable? The Privatisation of Women Abuse by Informal Networks in South Africa. *Journal of Gender and Religion in Africa*. 2012;(18)2:143-159.
6. Bougard, NB, Booyens, K. Adult Female Rape Victims' Views about the Thuthuzela Care Centres: A South African Multidisciplinary Service Delivery Model. *Acta Criminologica: Southern African Journal of Criminology*. 2015; 5: 19-33.
7. Johnson, S, Mahlalela, NB, Mills E. Client experience of rape victims accessing governmental post-rape services in South Africa. Brighton: Institute of Developmental Studies, 2017. [https://opendocs.ids.ac.uk/opendocs/bitstream/handle/20.500.12413/12857/Client%20experience\\_online.pdf](https://opendocs.ids.ac.uk/opendocs/bitstream/handle/20.500.12413/12857/Client%20experience_online.pdf) (accessed 14 July 2020).
8. The Foundation for Professional Development. Thuthuzela Care Centres Compliance Audit and Gap Analysis. Pretoria: The Foundation for Professional Development, 2016. <http://shukumisa.org.za/wp-content/uploads/2018/02/PA00MQJ6-1.pdf> (accessed 28 May 2020).
9. Creative Consulting and Development Works. Process Evaluation of NGO Services at Thuthuzela Care Centres. Cape Town: Networking HIV & AIDS Community of Southern Africa, 2018. <http://www.nacosa.org.za/wp-content/uploads/2018/09/GBV-Evaluation-Report-Web.pdf> (accessed 28 May 2020).
10. Vetten L. "It sucks/It's a wonderful service": Post-rape care and the micro-politics of institutions. Johannesburg: Shukumisa Campaign and ActionAid South Africa. <http://shukumisa.org.za/wp-content/uploads/2017/09/Thuthuzela-Care-Centres-Shukumisa-Report-2015.pdf> (accessed 28 May 2020).
11. National Shelter Movement of South Africa. Who we are & what we do. 2020. <https://www.nsmsa.org.za> (accessed 14 July 2020).
12. Watson J, Lopes C. Shelter Services to Domestic Violence Victims – Policy Approaches to Strengthening State Responses. Cape Town: Heinrich Böll Foundation, 2017. [https://za.boell.org/sites/default/files/policy\\_brief\\_final\\_02\\_web.pdf](https://za.boell.org/sites/default/files/policy_brief_final_02_web.pdf) (accessed 28 May 2020).
13. Commission for Gender Equality. Investigative Report: State of Shelters in SA. Braamfontein: Commission for Gender Equality, 2017. <http://www.cge.org.za/wp-content/uploads/2014/05/State-of-Shelters-in-SA.pdf> (accessed 28 May 2020).
14. Van der Heijden, I, Harries, J, Abrahams Naeemah. Barriers to gender-based violence services and support for women with disabilities in Cape Town, South Africa. *Disability & Society*. 2019.
15. Naidu, E, Mkhize N. Gender-Based Violence: The Lesbian and Gay Experience. Agenda: Empowering Women for Gender Equity. 2005;66(1):34-38.
16. United Nations Women. Violence against women and girls: the shadow pandemic. 2020. <https://www.unwomen.org/en/news/stories/2020/4/statement-ed-phumzile-violence-against-women-during-pandemic> (accessed 28 May 2020).
17. Onyango MA, Resnick K, Davis A, Shah RR. Gender Based Violence Among Adolescent Girls and Young Women: A Neglected Consequence of the West African Ebola Outbreak. In: Schwartz D., Anoko J., Abramowitz S. (eds) *Pregnant in the Time of Ebola. Global Maternal and Child Health (Medical, Anthropological, and Public Health Perspectives)*. Springer, Cham, 2019: 121-132. [https://doi.org/10.1007/978-3-319-97637-2\\_8](https://doi.org/10.1007/978-3-319-97637-2_8)
18. John N, Casey SE, Carino G, McGovern T. Lessons Never Learned: Crisis and gender-based violence. *Developing World Bioeth*. 2020; 00: 1– 4. <https://doi.org/10.1111/dewb.12261>
19. National Shelter Movement of South Africa. Domestic violence safety planning during the time of COVID-19. [https://za.boell.org/sites/default/files/2020-04/DVSafetyPlan\\_digital.pdf](https://za.boell.org/sites/default/files/2020-04/DVSafetyPlan_digital.pdf) (accessed 28 May 2020).
20. Adonis, N, Lopes C. GBV victims at risk & shelters, COVID-19 lockdown. Cape Town: Heinrich Boll Stiftung, 2020. <https://za.boell.org/en/2020/03/26/victims-gbv-are-some-those-most-risk-not-only-coronavirus-during-nationwide-lockdown> (accessed 14 July 2020).
21. Nicolson, G. Lockdown: Many 'trapped in their home with their abuser'. *Daily Maverick*, 2020. <https://www.dailymaverick.co.za/article/2020-04-08-lockdown-many-trapped-in-their-home-with-their-abuser/> (accessed 28 May 2020).
22. National Shelter Movement of South Africa. Press Release: GBV Victims & Shelters, COVID-19 Lockdown. 2020. <https://www.nsmsa.org.za/2020/03/26/hello-world/> (accessed 28 May 2020).
23. International Rescue Committee. Press Release: New Data Shows a Decrease in Women Being Able to Report Incidents of Domestic Violence in Fragile and Conflict-Affected Countries. 2020. <https://www.rescue.org/press-release/new-data-shows-decrease-women-being-able-report-incidents-domestic-violence-fragile> (accessed 14 July 2020).
24. Sibanyoni M. Infographic: South Africa's lockdown level 5, 4, 3, 2, and 1. SABC News, 2020. <https://www.sabcnews.com/sabcnews/infographic-south-africas-lockdown-level-5432-and-1/> (accessed 28 May 2020).
25. National Shelter Movement of South Africa. Shelter Directory. 2020. <https://www.nsmsa.org.za/shelter-directory/> (accessed 29 April 2020).
26. Department of Justice and Constitutional Development. Thuthuzela Care Centres (TCCs) Site Staff and Contact Details. <https://www.justice.gov.za/vg/TCCs-list.pdf> (accessed 29 April 2020).
27. Ntseku, M. Shelters taking strain as domestic violence rises during Covid-19 lockdown. IOL, 2020. <https://www.iol.co.za/capeargus/news/shelters-taking-strain-as-domestic-violence-rises-during-covid-19-lockdown-46683482> (accessed 14 July 2020).
28. Barnett, M, DC Grabowski. Nursing Homes Are Ground Zero for COVID-19 Pandemic. *JAMA Health Forum*. 2020.
29. Tobolowsky FA, Gonzales E, Self JL, et al. COVID-19 Outbreak Among Three Affiliated Homeless Service Sites - King County, Washington, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(17):523-526.
30. Breiding MJ, Basile KC, Klevens J, Smith SG. Economic Insecurity and Intimate Partner and Sexual Violence Victimization. *Am J Prev Med*. 2017;53(4):457-464.
31. Hatcher AM, Stöckl H, McBride R-S, Khumalo M, Christofides N. Pathways from food insecurity to intimate partner violence perpetration among peri-urban men in South Africa. *Am J Prev Med*. 2019;56(5):765-72.
32. Ramaphosa, C. Full text: Cyril Ramaphosa's speech on lockdown measures. Bulawayo 24 News. 2020. <https://bulawayo24.com/index-id-opinion-sc-speeches-byo-185301.html> (accessed 28 May 2020).